

CLIMATE SENSITIVITY AND THE VALUE OF AGRICULTURAL PRODUCTION IN THE BRAZILIAN NORTHEAST: AN APPROACH USING SPATIAL PANEL DATA

Helson Gomes de Souza* Edward Martins Costa†
Francisco José Silva Tabosa‡ Pablo Urano de Carvalho Castelar§

ABSTRACT

The present work aims to verify the sensitivity of Northeast agricultural production in relation to the changes in precipitation and temperature levels in the Northeast region of Brazil. For that, data from 2006 to 2016 were used for 952 municipalities in the Northeast region, inserted in an econometric methodology for panel data with spatial specification. Through this procedure it was concluded that in the last years the agricultural production of the Northeast of Brazil has become more sensitive to the changes in precipitation levels. In addition, it was concluded that public policies to combat the effects of climate change on agriculture in the Northeast region will have greater impacts when based on the water needs of municipalities.

Key words: climate Change. Agricultural Production. Environmental Economics.

RESUMO

O presente trabalho tem como objetivo verificar a sensibilidade da produção agrícola nordestina em relação às mudanças nos níveis de precipitação e temperatura na região Nordeste do Brasil. Para tanto, foram utilizados dados de 2006 a 2016 para 952 municípios da região Nordeste, inseridos em uma metodologia econométrica para dados em painel com especificação espacial. Através deste procedimento, concluiu-se que nos últimos anos a produção agrícola do Nordeste do Brasil se tornou mais sensível às mudanças nos níveis de precipitação. Além disso, concluiu-se que as políticas públicas para combater os efeitos das mudanças climáticas na agricultura da região Nordeste terão maiores impactos quando baseadas nas necessidades hídricas dos municípios.

Key words: Mudanças Climáticas. Produção Agrícola. Economia Ambiental.

Classificação JEL: Q10. Q54. Q50.

Área 2: Desenvolvimento Regional.

*Mestre em Economia Rural (MAER-UFC); Doutorando em Economia Aplicada (PPGE-UFPB). E-mail: helson.g.souza@gmail.com

†Professor do Departamento de Economia Agrícola da Universidade Federal do Ceará – UFC. E-mail: edward-costa@ufc.br

‡Professor do Departamento de Economia Agrícola da Universidade Federal do Ceará – UFC. E-mail: franzetabosa@ufc.br

§Professor do curso de Finanças da Universidade Federal do Ceará - UFC. E-mail: pcastelar@ufc.br

1 INTRODUCTION

It is reasonable to assume that climate conditions affect economic outcomes. The geographic scenario in which a country or region is located, and changes which occur therein, will naturally lead to different results in terms of production. Therefore, how these changes in climate conditions impact production, and by how much, has been cause for analysis and concern in recent years. The environmental impacts caused by climate change have been an issue that has gained relevance in the socioeconomic context since the late 20th and the beginning of the 21st century. In light of the recent effects of climate change on the environment and the global economy, tools have been put in place to mitigate the impacts of climate-related economic activities worldwide (KUNWAR; BOHARA, 2017).

Thus, nations have recently taken greater responsibility for the effects of climate change. For example, the formulation of global goals and international agreements aims at minimizing these effects. An example of this was the Paris Agreement, set up at the 21st Conference of Parties (COP21), in Paris, France. The agreement seeks to strengthen the global response to the threat of climate change and strengthen the capacity that countries have to deal with the impacts of climate change (Ministério do Meio Ambiente - Brazilian Ministry of Environment), (MMA, 2017).

Data from the National Aeronautics and Space Administration (NASA, 2017) show that the average global temperature has increased since the late nineteenth and early twentieth century, and that global temperature growth rates began to increase after the 1940s, and intensified after the 1970s.

Yalew et al. (2017) using a microeconomic approach, find that changes in the climate cause different socioeconomic impacts, that can be intensified depending on the sector analyzed. However, the authors demonstrate that the sector that is most impacted by climate change is agriculture. As a result, these impacts are more intense on economies which are mainly focused on the primary sector.

Nelson et al. (2009a) indicate that climate change will lead to future effects that will have major impacts, especially in economies based on the agricultural export sector. For the authors, underdeveloped countries located along and south of the Equator will be more affected. In this context, Rosegrant et al. (2008) forecast the influence of climate change and indicate a large future loss of production in Brazil, especially in relation to sugarcane, rice and soybean crops.

Filho et al. (2016) highlight the effects of climate change in some specific regions of Brazil. The authors verified that the changes in the climate make it difficult to exit the poverty condition in rural areas in the North and Northeast regions of the country. In addition, the agricultural sector of the Northeast of Brazil suffers greater impacts from the changes in the climate behavior, especially when it comes to family and subsistence farming.

In this context, the Brazilian National Institute for Space Research (INPE, 2015) reports that the general trend indicates a reduction in the level of precipitation in the Brazilian Northeast, as well as the temperature diverging in relation to its historical average in the region. For Filho et al. (2016), the main consequences of this phenomenon are the loss of profitability of agriculture in the region and the intensification of poverty and social precariousness in the rural areas of the Northeast.

As discussed by Nunes (2016), the different regions of a country, particularly one of continental dimensions such as Brazil, suffer disparate effects because of climate change. In particular, says the author, the North and Northeast regions are the most vulnerable to the current shifts in the climate, because these regions are considered extremes in terms of environment. The author bases her statement on the Climate Report of the Ministry of the Environment, the Secretariat of Biodiversity and Forests and the Directorate of Biodiversity Conservation of 2015. Obermaier e Rosa (2013), likewise, also point out that the semi-arid regions, such as the Brazilian Northeast, tend to feel a greater impact of the variations in the climate.

In view of these considerations, the present work seeks to answer the question: What are the impacts of climate change on the agricultural production in the Northeast of Brazil? Therefore, the main objective of this research is to verify the sensitivity of the Brazilian Northeast agricultural

production in relation to the changes in precipitation and temperature levels. The hypothesis is that the agricultural production of the Brazilian Northeast is spatially autocorrelated, generating the spatial overflow of the changes in the production of a certain region to the neighboring spaces, or municipalities.

The contribution intended concerns the quantification of the sensitivity that the agricultural sector has to climate change, in a region where the economy is highly dependent on agriculture. The present work is subdivided into five sessions, including this brief introduction. The second, which follows, encompasses the theoretical and literary foundation under which the work is based. The third refers to the methodological framework used. The fourth discusses the results found. Lastly, the concluding remarks are presented.

2 LITERATURE REVIEW

2.1 Climate change and its Effects on the Agricultural Sector

Climate change has had worrying effects in recent years, and these shifts have a stronger impact on the agricultural sector, as discussed by various authors. This section intends to highlight only a few of those discussions, in order to motivate the analysis performed.

In this sense, the economic literature has been giving significant prominence to works that analyze the behavior of agriculture in relation to climatic variations. Note, for example, [Piao et al. \(2010\)](#), [Parry et al. \(2004\)](#), [Fischer, Shah e Velthuizen \(2002\)](#), and [Wiebe et al. \(2015\)](#), all of which, on some level, find correlation between shifts in the climate and economic productivity.

According to [Nelson et al. \(2009b\)](#), climate change has effects on the agricultural sector and affects other sectors through existing sectoral integrations. Changes in the levels of greenhouse gas emissions, temperature and precipitation cause direct changes in agricultural production, which generate impacts on the prices of primary extraction products, impacting, thus, other sectors. [Parry et al. \(2004\)](#) forecast global food production, taking into account the effect of climate change and the characteristics of population growth in the world in recent years. The authors conclude that the consequences of climate change can be catastrophic over the next 60 years. For these authors, it is necessary to implement long-term policies aimed at preventing the impacts suffered by the agricultural sector caused by climate change, such as reducing emissions of carbon dioxide (CO_2) and other greenhouse gases.

In the view of [Calzadilla et al. \(2013\)](#), the greatest impact on the agricultural sector as a result of climate change is on the productivity of the factors of production. While analyzing assesses the potential impacts of climate change and CO_2 fertilization on global agriculture, using a version of the GTAP-W model, which distinguishes between rainfed and irrigated agriculture and implements water as an explicit factor of production for irrigated agriculture, the authors predict results for agricultural production through projections up to the year 2050, taking into account the effects of climate change on the level of production of agriculture and the well-being of individuals. [Calzadilla et al. \(2013\)](#) conclude that if the behavior of climate variables continues at the current pace, agricultural production and welfare will fall worldwide in the next three decades.

Using a different methodological approach, [Evangelista et al. \(2013\)](#) determine that the analyses of the impact of changes in the climatic variables on the agricultural sector should consider the spatial factor. The authors consider the modifications in the levels of precipitation and temperature and their impacts on the agricultural production in Ethiopia, where nearly all of the country's agriculture is dependent on rainfall. [Evangelista, Young e Burnett \(2013\)](#) verify, then, using a Maxent software fit with crop data collected from household surveys and bioclimatic variables from the WorldClim database to develop spatially explicit models of crop production in Ethiopia, that these changes exhibit a common spatial behavior during the year. In addition, the authors determine that the impacts of climate change on agriculture are spatially concentrated in some periods and

subsequently spread throughout the area under analysis.

Ju et al. (2013), the changes in the temperature levels of the last decades have caused considerable impacts on world agricultural production, which requires the development of adaptation methods to these changes. However, the authors indicate that hydric changes resulting from the alteration of rainfall levels have a greater impact on the agricultural sector. This result is also verified by Kunwar e Bohara (2017), where the authors observe that the values of rural properties show sensitivity to the changes of precipitations superior to the sensitivity of the changes in the temperature.

The approach developed by Ochieng, Kirimi e Mathenge (2016), who estimate the effects of climate variability and change on agricultural revenues of small-scale farmers in Kenya, also indicates that agricultural production can be impacted by changes in temperature and precipitation levels. The authors, using a balanced panel household data set collected in 2000, 2004, 2007 and 2010 from eight agro-regional zones and climate data from 1980 to 2010, argue that the interactions between these two variables directly affect the behavior of agriculture. In addition, the authors indicate an increase in agricultural production elasticities related to precipitation and temperature levels in a projection for the next three decades.

Performing a literary survey on climate issues in Brazil, Moraes e Filho (2013) point out that it is necessary to understand the knowledge points concerning the issue of climate change in Brazil, at least in its economic aspects. Furthermore, the authors argue that only from having such knowledge it is possible to think about strategies, being this the main contribution of the works in the area. Also, they emphasize that, in Brazil, mitigation policies are superior, in economic terms, to the perspectives of vulnerability and/or adaptation to climate issues. However, in their opinion, more consistent information is still necessary for society to plan its lines of action.

Cunha et al. (2013) emphasize that climate change in Brazil affects mainly small producers. Using a treatment effect model with propensity score matching, the authors argue that rainfed family agriculture is the most affected by these phenomena, thus this type of production has a high sensitivity with regard to changes in precipitation levels. Therefore, the authors stress that it is necessary to implement measures that facilitate the technological adaptation of small rural producers to climate variability in Brazil. Also, Cunha et al. (2013) cite irrigation as an adaptive measure to balance the effects of climate change on agriculture and highlight the importance of policies to support the expansion of this technique for small farmers.

In this sense, Araújo et al. (2014) analyze the effect of climate change on the agricultural productivity levels of the states of Northeast Brazil, considering the impact of temperature and precipitation levels on the productivity of crops such as maize, sugar cane and manioc. The authors, using a tobit model for panel data, conclude that the productivity levels of the crops studied could be much higher than those that would be observable if there were no changes in the climate. In addition, states such as Rio Grande do Norte, Paraíba and Pernambuco may show productivity losses in the three analyzed crops, both in the medium and long run. The authors also suggest that climate change will negatively impact productivity levels in the municipalities of the South and Center-South of the state of Bahia.

From another perspective, Domingues, Magalhães e Ruiz (2011), analyze the impact of climate change in the Brazilian Northeast from estimates of the implications on the availability of suitable land for agricultural activity in a set of crops. The authors, using computable general equilibrium models, conclude that climate change increases the potential for economic losses in the Northeast region of the country, especially in the poorest states, pointing to the need for mitigation and emission control policies. These authors further indicate that, if measures to prevent and lessen the effects of climate change are not implemented, the economic effects on employment, for example, could have a significant impact on migratory flows, impacting the high pressure on urban infrastructure of the various metropolises of the Northeast and other regions of the country.

According to the work developed by Filho et al. (2016). agriculture in the Brazilian North-

east presents a high vulnerability to the changes in the climate, mainly in what concerns shifts in pluviometry. The authors point out that in the coming years, the effects of climate change on agriculture in the Northeast of Brazil will be intensified in a higher proportion than the other regions, and hydric changes are the main issues responsible for the problem. The authors also point out that family farming in the Northeastern region will suffer greater impacts from climate change than other subsectors, thus arguing for the need to implement policies aimed at controlling and supplying water to the region.

3 METHODOLOGY

3.1 Initial Specification

While analyzing the impact of climate change on agriculture, [Kunwar e Bohara \(2017\)](#) assume that the value of agricultural land is impacted by variations in rainfall and local temperature. Thus, the authors use a spatial methodology for panel data to measure the impact of climate change on the value of rural properties. Considering a Ricardian approach, [Kunwar e Bohara \(2017\)](#) assume that the value of rural properties is a function of the value of the average precipitation, average temperature, the deviations in precipitation and temperature, and the squared values of those variables.

In the present work, the approach used by [Kunwar e Bohara \(2017\)](#) is applied to the production value of municipalities in the Brazilian Northeast. However, it is assumed here that the production value is also impacted by the interactions between precipitation and temperature. In this sense, time and space data are used, applied to a spatial econometric methodology. The following subsections are intended to present the tools used for this procedure.

3.2 Data

The data used here comes from two different sources. In relation to the value of agricultural production, the sum of the total value of the production of temporary and permanent crops is reported in the Municipal Agricultural Production ([PAM, 2017](#)), which is made available annually by the Brazilian Institute of Geography and Statistics ([IBGE, 2017](#)) in the IBGE System of Automatic Recovery ([SIDRA, 2017](#)). The value of production was corrected for December 2016 values by the Brazilian Extended Consumer Price Index (IPCA).

Regarding the climate variables, the information was obtained in the Agrometeorological Monitoring System ([AGRITEMPO, 2017](#)). The mean temperature and precipitation were constructed using the arithmetic mean of the maximum and minimum daily estimates for each variable.

For the construction of climate variables by seasons, the arithmetic mean of the daily mean values of precipitation and temperature was used. Following the specifications of INPE, the period between December 21 and March 19 was considered as summer; March 20 to June 20 as autumn; June 21 to September 21 as winter and September 22 to December 20 as spring¹.

After the homogenization of the data, a total of 952 municipalities² of the Brazilian Northeast region comprise the database, being 30 municipalities of the state of Sergipe, 37 municipalities in the state of Alagoas, 290 in the state of Bahia, 83 in the state of Pernambuco, 67 in the state of Paraíba, 55 in the state of Rio Grande do Norte, 112 in Ceará, 338 in Piauí, and 140 in the state of Maranhão. Thus, the *QGIS* software was used to construct a digital layer in the shapefile format

¹It should be noted that in much of the Brazilian Northeast, climate seasons are not temporally well defined as specified, however, given the great variability in the definition of the seasons, and the lack of concrete information about the time in which these seasons occur in each municipality of the region, the authors decided to consider the period of the four conventional climate seasons as defined by INPE.

²The other municipalities were disregarded in view of the lack of information on climate variables or the value of agricultural production.

using points for the delimitation of the geographic location of each municipality, and afterwards, the *Geoda* software was used for the construction of the spatial weights matrices and calculation of the spatial autocorrelation indicators.

The period between 2006 and 2016 was chosen for this work. This choice was due to the greater availability of information on the climate variables featured in the *Agritempo* system.

3.3 Spatial Proximity Matrix

As the present work uses data distributed in time and space, the first step to be considered is to model the proximity of the spatial units in a numerical way. For this purpose, a spatial proximity matrix is constructed, consisting of an instrument capable of identifying neighboring units of a given region by means of the representation of areas in numerical terms. Based on the specification given by Almeida (2012), the neighborhood matrix has the following structure:

$$W_{ij} = \begin{cases} 1, & \text{if } i \text{ and } j \text{ are neighbors} \\ 0, & \text{if } i \text{ and } j \text{ are not neighbors} \end{cases} \quad (1)$$

3.4 Global Spatial Autocorrelation

When dealing with spatial data, before performing any econometric procedure, it is necessary to make an analysis of the characteristics of the spatial distribution of the data. This analysis is done in the present work through the verification of spatial autocorrelation, using the Moran Global Index, presented by Almeida (2012) as:

$$I = \frac{\sum_i \sum_j W_{ij} (Z_i - \bar{Z})(Z_j - \bar{Z})}{\sum_i^n (Z_i - \bar{Z})^2} \quad (2)$$

Where n represents the number of regions analyzed, Z_i is the value of the analyzed variable in area i , \bar{Z} is the mean value of the variable in a certain area, and W_{ij} represents the values indicated by the spatial proximity matrix used. The rejection of the null hypothesis of Moran's I , together with the positive sign, indicates the presence of spatial autocorrelation in the data. Non-rejection of the null hypothesis or the negative sign indicates a random dispersion of the data.

3.5 Econometric Methodology

In order to verify the sensitivity of agricultural production to climate change, the present work makes use of the method proposed by Kunwar e Bohara (2017), which uses a Ricardian approach to measure the sensitivity of the value of rural establishments to climate change in Nepal.

Initially, a conventional model for panel data is considered, which is given by:

$$y_{it} = \alpha_i + \beta Cmed_{it} + \theta Cmed_{it}^2 + \rho Cestdv_{it} + \nu Cest_{it}^2 + \omega (prec_{it} * temp_{it}) + \Gamma (prec_{it} * temp_{it})^2 + \phi control + \epsilon_{it} \quad (3)$$

Where $\alpha_1, \dots, \alpha_n$ is a vector corresponding to the fixed effects. y is the natural logarithm of the value of agricultural production, $Cmed_{it}$ is the annual mean of the climate variable (temperature and precipitation)³, $Cestdv_{it}$ represents the deviations of the climatic variables by season of the year, $prec$ represents the precipitation, $temp$ is the temperature and ϵ_{it} is the random error.

³Considering y as the value of rural establishments and X as the explanatory variables. Kunwar and Bohara (2017) consider a model such that $y = f(X, X_2, X_{dv})$. However, considering the problem of collinearity with the data used, the climate variables by season were only approached in the present work in its quadratic form.

However, [Anselin \(1999\)](#) indicates that when dealing with spatial data, the assumption of homogeneity must be addressed in the econometric procedures used. Non-homogeneity may indicate that the coefficient of an explanatory variable is statistically significant in a given municipality and not significant in another spatial unit.

Some techniques for crossection estimations are able to control this problem. However, these operations may not produce the same effects for stacked data. Thus, [Elhorst \(2003\)](#) indicates that the combination of stacked data with spatial models is one of the most robust tools in the field of spatial econometrics and that there are still unanswered questions about these methods, such as the control of spatial heterogeneity over the estimated coefficients.

[Barreto, Almeida e Lima \(2010\)](#) use fixed effects to control spatial heterogeneity, a procedure also adopted in the present investigation. Still, [Baylis, Paulson e Piras \(2011\)](#) emphasize that in this type of approach, one must also reduce errors by including factors external to the climatic variables that affect agriculture in the different units of space such as population, infrastructure, income, demographic density, conditioning factors irrigation etc. Therefore, the reduction of the effects from stochastic errors can be achieved by inserting a set of control variables into the estimates⁴.

In the present work, control variables were incorporated in view of data availability and taking into account the procedures used in literature as done by [Baylis, Paulson e Piras \(2011\)](#) and [Kunwar e Bohara \(2017\)](#). With this, the natural logarithm of the Municipal Gross Domestic Product (GDP)⁵, the average area available per inhabitant⁶ and the natural logarithm of the municipal population⁷.

Beginning from Equation 4 and using the procedure of incorporation of the spatial effects given by [Elhorst \(2010\)](#), one arrives at a general fixed effects model that includes the spatial dependency including spatial lags to have control of the spatial autocorrelation, being given by:

$$\begin{aligned}
y_{it} &= \alpha_i + \rho W_1 y_{it} + \beta C med_{it} + \theta C med_{it}^2 + \varrho C est dv_{it} + \nu C est_{it}^2 + \omega (prec_{it} * temp_{it}) \\
&+ \Gamma (prec_{it} * temp_{it})^2 + \phi control + W_1 (\beta C med_{it} + \theta C med_{it}^2 \\
&+ \varrho C est dv_{it} + \nu C est_{it}^2 + \omega (prec_{it} * temp_{it})) \\
&+ \Gamma (prec_{it} * temp_{it})^2 + \phi control) \tau + \xi_t \\
\text{with } \xi_t &= \lambda W_2 \xi_t + \epsilon_t
\end{aligned} \tag{4}$$

Where $W_1 y_t$ is the dependent variable spatially lagged, $W_2 \epsilon_t$ are the error terms spatially lagged, W is the neighborhood matrix, and λ and ρ are the scalar spatial parameters, being τ a vector of spatial coefficients.

The general model of random effects with spatial dependence is presented by [Almeida \(2012\)](#) as:

$$\begin{aligned}
y_{it} &= \rho W y_{it} + \beta C med_{it} + \theta C med_{it}^2 + \varrho C est dv_{it} + \nu C est_{it}^2 + \omega (prec_{it} * temp_{it}) \\
&+ \Gamma (prec_{it} * temp_{it})^2 + \phi control + W (\beta C med_{it} + \theta C med_{it}^2 \\
&+ \varrho C est dv_{it} + \nu C est_{it}^2 + \omega (prec_{it} * temp_{it})) \\
&+ \Gamma (prec_{it} * temp_{it})^2 + \phi control) \tau + \xi_t \\
\text{with } \xi_t &= \lambda W \xi_t + \epsilon_t
\end{aligned} \tag{5}$$

⁴These variables were chosen based on various works in the literature, such as [Baylis, Paulson e Piras \(2011\)](#) and [Kunwar e Bohara \(2017\)](#).

⁵Made available annually by the IBGE. In view of the non-availability of this information for the year 2016, an estimate was considered, which was reached by multiplying the growth rate of the GDPs of 2014 and 2015 with the GDP of 2015.

⁶[Baylis, Paulson e Piras \(2011\)](#) use population density. However, considering the collinearity problem of this variable with the data used here, it was decided to use the average area per inhabitant, given by Area (municipal) / Population.

⁷Population estimates made available by the IBGE were used.

Imposing the restriction $\rho \neq 0$, $\tau = 0$ and $\lambda = 0$ to the model specified in Equation (4), one obtains the fixed effects model with spatial lags given by:

$$y_{it} = \alpha + \rho y_{it} + \beta Cmed_{it} + \theta Cmed_{it}^2 + \varrho Cestdv_{it} + \nu Cest_{it}^2 + \omega(prec_{it} * temp_{it}) + \Gamma(prec_{it} * temp_{it})^2 + \phi control + \epsilon_t \quad (6)$$

Imposing the restriction $\rho \neq 0$, $\tau = 0$ and $\lambda = 0$ to the model specified in Equation (5), one obtains the random effects model with spatial lags given by:

$$y_{it} = \rho W y_{it} + \beta Cmed_{it} + \theta Cmed_{it}^2 + \varrho Cestdv_{it} + \nu Cest_{it}^2 + \omega(prec_{it} * temp_{it}) + \Gamma(prec_{it} * temp_{it})^2 + \phi control + \xi_t$$

with $\xi_t = \alpha + \epsilon_t$

The spatial error model with fixed effects is also considered, which is characterized by featuring the spatial autocorrelation in the form of an autoregressive error. In order to obtain this model, the constraints $\rho = 0$, $\tau = 0$ and $\lambda \neq 0$ are imposed on Equation (4), to achieve:

$$y_{it} = \alpha + \beta Cmed_{it} + \theta Cmed_{it}^2 + \varrho Cestdv_{it} + \nu Cest_{it}^2 + \omega(prec_{it} * temp_{it}) + \Gamma(prec_{it} * temp_{it})^2 + \phi control + \xi_t$$

with $\xi_t = \lambda W \xi_t + \epsilon_t$

For the spatial error model with random effects, the constraints $\rho = 0$, $\tau = 0$ and $\lambda \neq 0$ in Equation (5) yield:

$$y_{it} = \beta Cmed_{it} + \theta Cmed_{it}^2 + \varrho Cestdv_{it} + \nu Cest_{it}^2 + \omega(prec_{it} * temp_{it}) + \Gamma(prec_{it} * temp_{it})^2 + \phi control + \xi_t$$

with $\xi_t = \alpha + \lambda W \xi_t + \epsilon_t$

The choice of the more appropriate model among the estimates is made using the criteria indicated by Almeida [Almeida \(2012\)](#), where the choice of the appropriate model should be made according to: (i) verifying if the observed effects should be included in the estimates; (ii) choosing the most appropriate model with unobserved effects; (iii) the appropriate model with no observed effects is estimated; (iv) checking the spatial dependence of the residue; (v) verifying the presence of spatial autocorrelation in the residues of the model with unobserved effects, one estimates the panel data model encompassing spatial autocorrelation; (vi) the model that does not present spatial autocorrelation in the residues and that presents the least information criterion is chosen.

If the model chosen is the spatial lag model, as shown in Equation (6), it is possible to obtain the local and total impacts of the climate variations by calculating the marginal effects, which are given by [LeSage e Pace \(2009\)](#) as:

$$\frac{\delta y_i}{\delta x_{ir}} = S_r(W)_{ii}$$

and

$$\frac{\delta y_i}{\delta x_{jr}} = S_r(W)_{ij} \quad (10)$$

In which $S_r(W) = (I_n - \phi W)^{-1} \beta_r$, i and j represent municipalities i and j respectively. β_r is the coefficient of the r -th variable and $S_r(W)$ is an $(n \times n)$ matrix where the elements contained in the main diagonal represent the direct impacts and the elements outside the main diagonal represent the indirect impacts. The sum of these two impacts forms the total impact. Once the estimates have been made and the appropriate model chosen, the next step is to calculate the sensitivity of the agricultural production in relation to the level of precipitation and the temperature. Following the procedure of Kunwar and Bohara (2017), and including the interactions between precipitation and temperature, this process is given by:

$$\begin{aligned}
 E\left(\frac{\delta \ln(y)}{\delta \text{prec}}\right) &= \beta + 2 * \theta + \omega * \text{temp} + 2 * \Gamma * \text{prec} * \text{temp}^2 \\
 &\text{and} \\
 E\left(\frac{\delta \ln(y)}{\delta \text{temp}}\right) &= \beta + 2 * \theta + \omega * \text{temp} + 2 * \Gamma * \text{prec}^2 * \text{temp}
 \end{aligned}
 \tag{11}$$

4 RESULTS

To specify the characteristics of the data used, Table 1 shows the descriptive statistics of the variables. Mean precipitation and mean temperature were obtained by the ratio of the sum of the values and the number of days of the period⁸.

⁸Annual and seasonal averages of daily values of temperature and precipitation were used, that is: $\frac{\sum(\text{temp} * \text{prec})_{\text{day}}}{\sum(\text{days})}$

Table 1: Descriptive statistics of the variables

Variable	Observations	Mean	Standard Deviation
Ln(production)	10.472	8.5658	1.8538
Precipitation	10.742	2.7175	1.7194
Temperature	10.742	25.7391	1.7194
<i>Precipitation</i> ²	10.742	10.3410	24.0820
<i>Temperature</i> ²	10.742	669.5832	128.0241
prec*temp	10.742	69.5731	40.8843
(<i>prec * temp</i>) ²	10.742	6511.7760	9036.4000
<i>Prec_spring</i> ²	10.742	7.2106	24.8750
<i>Prec_summer</i> ²	10.742	27.3372	40.5980
<i>Prec_autumn</i> ²	10.742	29.4538	51.0579
<i>Prec_winter</i> ²	10.742	1.9986	21.9598
<i>Temp_spring</i> ²	10.742	728.3846	136.2860
<i>Temp_summer</i> ²	10.742	711.5912	111.5421
<i>Temp_autumn</i> ²	10.742	646.8529	129.8654
<i>Temp_winter</i> ²	10.742	599.9043	155.3370
Temp_spring_dv	10.742	1.1144	0.8214
Temp_summer_dv	10.742	0.8417	1.1459
Temp_autumn_dv	10.742	-0.4642	0.7455
Temp_winter_dv	10.742	1.4836	1.1371
Prec_spring_dv	10.742	-0.8449	1.9794
Prec_summer_dv	10.742	1.5985	1.7991
Prec_autumn_dv	10.742	1.4435	2.2507
Prec_winter_dv	10.742	-2.1353	1.5046
Control			
Area/Population	10.742	0.0785	0.1192
LnGDP_percapita	10.742	8.9505	0.5953
Lnpopulation	10.742	9.7994	0.9722

Source: Elaborated by the authors based on estimations from [AGRITEMPO \(2017\)](#).

Note: *Pre* = precipitation (mm); *temp* = temperature (°c); *dv* = standard deviation.

The average precipitation of the analyzed period was 2.7175 mm. which, in accumulated values, would generate an average annual precipitation of approximately 990 mm. It is also observed that the greatest precipitations occur during the summer and fall, which corresponds to the period from December 21 to June 20.

The average temperature for the analyzed period was 25.7391°C. Checking the square of the averages by seasons, it is observed that the highest temperatures occur in the spring, that is, between September 22 and December 20.

The standard deviations inform how much the climate variables distance themselves from the average, allowing an approximation of the quantification of the climate changes. Regarding precipitation and temperature, Table 1 shows that, on average, the greatest climate changes occur during the winter, that is, between June 21 to September 21.

Before making any statistical inference, it is necessary to verify the temporal characteristics of the data. Therefore, the stationarity condition of the variables was analyzed by the Levin-Lin-Chu stationarity test. The results of this procedure are shown in Table 9, featured in the appendix. The rejection of the null hypothesis confirms that all the variables used are distributed around a fixed mean, with constant variance over time, which indicates the stationarity condition of the series.

Table 2 presents the results obtained with the Moran Global Index for the agricultural production of the analyzed municipalities. For all the years considered in the sample, the Moran Global Index presented a positive value and was statistically significant. This result indicates that the agricultural production of the studied municipalities is spatially autocorrelated. This implies a spatial overflow of production, that is, changes occurred in the agricultural production of a given municipality overflow in spatial terms, causing changes in the agricultural production of the neighboring municipalities.

Table 2: Global spatial autocorrelation of the value of agricultural production.

	Moran's I	P - Value
2006	0.477	0.0001
2007	0.513	0.0001
2008	0.498	0.0000
2009	0.482	0.0010
2010	0.555	0.0000
2011	0.494	0.0002
2012	0.474	0.0000
2013	0.524	0.0003
2014	0.506	0.0000
2015	0.522	0.0001
2016	0.458	0.0000

Source: Elaborated by the authors.

Note: The statistical significance considered for this work is the 5% level.

After verifying the presence of spatial autocorrelation in the data used, the next step is to measure the sensitivity of agricultural production in relation to precipitation and temperature by means of a procedure that encompasses the effects stemming from these phenomena. In the present work this process is done through a spatial data panel. In order to decide which estimator should be considered, an F test was initially used to verify the possibility of estimation of the models through Ordinary Least Squares via a pooled model. With this procedure, the null hypothesis of there being no significant effects was rejected. In addition, a Breusch-Pagan test was performed and it was verified that it is not possible to ignore the presence of significant effects, which can only be adequately treated through the Fixed Effects estimator.

Once the estimates are obtained, one must check whether the effects should be treated as fixed or random. For that purpose, the Hausman test is used in this work. It is verified that in all the estimations the random effects are invalid, and, therefore, the fixed effects model must be considered to answer the problem here. The results of these procedures are presented in Table 3.

For models with spatial specification, a neighborhood matrix of type K neighbors with $K = 1$ was used. The choice of this matrix was determined by the criteria indicated by [Baumont \(2004\)](#). This mechanism consists in estimating the model without the spatial effects, obtaining the model residues and verifying the spatial autocorrelation of the residues through the Moran Global Index, taking into account a set of neighborhood matrices⁹. Then, the matrix chosen will be the one that obtains the greatest statistically significant value for the Moran Global Index applied to the residuals of the model without spatial effects.

In order to choose the model that explains the problem in question more adequately, the procedure indicated by [Almeida \(2012\)](#) was used. as specified in subsection 3.4. Having fulfilled the prepositions (i), (ii), (iii), (iv) and (v), the absence of spatial autocorrelation in the spatial specification of the fixed effects models was verified. Thus, the appropriate model that explains the

⁹The results for this procedure are described in the appendix. Table 8.

relationships analyzed is the spatial lag model, considering that it presented the lowest information criterion.

Analyzing the spatial lag model, it is verified that the spatial autocorrelation coefficient (ρ) features a positive signal and was statistically significant. This result may indicate the existence of clusters for agricultural production in the Northeast region of Brazil. Furthermore, the value obtained with the coefficient (ρ) indicates that the changes in the production of a given municipality overflow in spatial terms, also impacting neighboring municipalities. As a consequence of this result, it should be pointed out that a public policy aimed at impacting the agricultural production of the Northeast of Brazil should take into consideration the possibility of capturing the effects arising from spatial proximity.

The squares of the levels of precipitation by season show that increased precipitation in autumn reduces the value of agricultural production in the Brazilian Northeast. The values obtained with the coefficients of the temperatures per season of the year in their quadratic form indicate that temperature increases in autumn reduce agricultural production, while the increase in winter temperatures leads to greater values of agricultural production in the Northeast.

The results obtained with the temperature deviations show that the increase in the positive distance between the temperature in autumn and the average annual temperature causes increases in the value of agricultural production in the Northeast. In relation to precipitation deviations, it is observed that the increase in the positive distance between spring precipitation and the annual average precipitation reduces the production value in the region.

Regarding the control variables, the results show that the increase of the available area for each inhabitant as well as for the municipal population in general reduces the value of agricultural production.

Table 3: Estimation results using Fixed Effects.

	Normal	Lag	Error
Constant	8.3041*	-	-
ρ	-	0.2231*	-
λ	-	-	0.2137*
Prec	-0.7214*	-0.5836*	-0.6375
Temp	0.1829*	0.1506*	0.1422*
<i>Precipitation</i> ²	0.0253*	0.0202*	0.0209*
<i>Temperature</i> ²	0.0037	0.0029	0.0043
Prec*Temp	0.0411*	0.0332*	0.0366*
<i>(Prec * temp)</i> ²	0.0001*	0.0001*	0.0001*
<i>Prec_spring</i> ²	0.0010	0.0007	0.0005
<i>Prec_summer</i> ²	-0.0011	-0.0007	-0.0009
<i>Prec_autumn</i> ²	-0.0025*	-0.0018*	-0.0024*
<i>Prec_winter</i> ²	-0.0008	-0.0006	-0.0007
<i>Temp_spring</i> ²	-0.0003	-0.0007	-0.0008
<i>Temp_summer</i> ²	-0.0016	-0.0014	-0.0014
<i>Temp_autumn</i> ²	-0.0011*	-0.0086*	-0.0104*
<i>Temp_winter</i> ²	0.0030*	0.0026*	0.0029
Temp_spring_dv	0.0627	0.0687	0.0953
Temp_summer_dv	0.0682	0.0569	0.0658
Temp_autumn_dv	0.4266*	0.3305*	0.4025*
Temp_winter_dv	-0.0998	-0.0863	-0.0906
Prec_spring_dv	-0.0327*	-0.0258*	-0.0375*
Prec_summer_dv	0.0224	0.0158	0.0205
Prec_autumn_dv	0.0281*	0.0209	0.0328*
Prec_winter_dv	-0.0205	-0.0168	-0.0205
Control			
Area/Population	-1.5516*	-1.3985*	-1.2614*
Ln_GDP_Percapita	0.0109	0.0104	0.0115
Lnpopulation	-0.0948*	-0.0893*	-0.0781
Akaike		20171.55	20254.84

Source: Elaborated by the authors.

Note: Values succeeded by (*) represent statistical significance 5% level. Note: Prec = precipitation (mm); temp = temperature (°c); dv = standard deviation.

One of the advantages of estimating the spatial lag model is the possibility of quantifying spatial overflow through direct, indirect and total marginal effects¹⁰. The results for this procedure are presented in the appendix in Table 7. The averages for the marginal effects of the quadratic variables are outlined in Table 4. The average direct effects of the squares of the precipitation levels indicate that an increase in precipitation of a given municipality in 1 mm during the autumn season reduces the value of the agricultural production of that municipality by approximately 0.0155%. The indirect marginal effects of this variable indicate that the increase of 1 mm in the average autumn rainfall in a given municipality reduces by approximately 0.0038% the value of the agricultural production of neighboring municipalities. The overall impact of the 1mm increase in the average temperature in the fall is a reduction of 0.0194% in the value of agricultural production.

Regarding the squares of the temperatures by seasons, the direct effects show that the increase

¹⁰Since the variables by season of the year are squared, the marginal effects for these variables were calculated considering $Efeito(prec, temp)_{estação} = 2 * \nu * (prec, temp)_{estação}$ season.

of 1 mm in the average autumn and winter temperatures of a given municipality reduces by 0.4523% and increases by 0.1307%, respectively. the value of agricultural production in that municipality. On the other hand, the indirect effects indicate that the increase of 1 °c in the temperature of autumn and winter in a certain municipality overflows spatially, reducing by 0.1127% and 0.0325% the value of the agricultural production of the neighboring municipalities. Also, the total effects indicate that the 1 °c increase in autumn and winter temperatures in municipality i reduces by 0.5664% and increases by 0.1631%, respectively, the value of the agricultural production in that municipality and in the neighboring municipalities.

Table 4: Average marginal effects for climatic variables by season.

	Direct	Indirect	Total
<i>Prec_autumn</i> ²	0.0155	-0.0038	-0.0194
<i>Temp_autumn</i> ²	-0.4523	-0.1127	-0.5664
<i>Temp_winter</i> ²	0.1304	0.0325	0.1631

Source: Elaborated by the authors.

Continuing the analysis of the marginal effects of the fixed-effects model with spatial lags, it can be observed that the values of the direct effects for the quadratic temperature in the autumn and winter seasons and the deviations of the temperature in autumn and the precipitation in spring differ from the value of their respective coefficients in the estimation described in Table 3. According to [LeSage e Pace \(2009\)](#), this difference occurs as a result of the so-called feedback effect, which denotes changes in the value of agricultural production, which overflow in spatial terms and with the passage of time returns to the municipality of origin.

Regarding the deviations of temperatures per season, the direct effects indicate that the increase of 1 °c in the fall temperature in relation to the annual average temperature causes a local elevation of 0.3432% in agricultural production. On the other hand, the indirect effects indicate that the increase of 1 °c in the temperature of autumn in relation to the average annual temperature of a given municipality increases by 0.0853% the agricultural production of its neighbors. The total effects indicate that the overall impact of this change is 0.4285% on the value of agricultural production.

The direct marginal effects of seasonal precipitation deviations indicate that a 1 mm rise in the distance between the average precipitation in the spring and the average annual precipitation of a municipality reduces its agricultural production by 0.0259%, and in 0.0064% the production of neighboring municipalities, respectively. The total effects show that the overall impact of this change, considering the specific municipality and its neighbors, is a reduction of 0.0323% in the value of agricultural production.

The impacts of temperature and precipitation on agricultural production can only be visualized by calculating the sensitivities as denoted in Equation 11. This procedure is demonstrated in terms of averages for the Northeast region of Brazil in Figure 1.


From 2006 to 2011, agricultural production in the Brazilian Northeast was more sensitive to changes in temperature levels than to changes in average precipitation. From that point on, agricultural production became more sensitive to changes in precipitation levels, with the exception of 2014. Also, comparing the year 2016 with the beginning of the analyzed period, agricultural production in the Northeast region became more sensitive to changes in precipitation levels and less sensitive to changes in temperature.

The trajectory of the sensitivities outlined in Figure 1 demonstrates an already expected and well-known conclusion, which indicates a great dependence of the agricultural sector on rainfall stability and that public policies to combat the effects of climate change on agriculture in the Brazilian Northeast region will have greater impacts when based on hydric soil characteristics.

The high levels of sensitivity of the Northeastern agricultural production to climate change, as well as the intensification of the average sensitivity to the changes in pluviometry described in

Figure 1, expose producers to a situation already verified by [Burke e Emerick \(2016\)](#), which indicate that farmers need adaptation to the new climate scenarios, mainly with respect to technology and other factors that influence agricultural productivity.

Figure 1: Average sensitivity of agricultural production in relation to precipitation and temperature.



Source: Elaborated by the authors using the research data.

The intensification of agricultural production sensitivity in relation to changes in average precipitation levels from 2014 can be explained mainly by the worsening water scarcity that has been installed in most of the municipalities of the Brazilian Northeast in this period, as shown by the World Bank document organized by [De Nys, Engle e Magalhães \(2016\)](#).

Thus, it can be stated that the control of temperature shifts is an important mechanism for the adequate performance of agricultural production in the Brazilian Northeast. However, as stated by [Filho et al. \(2016\)](#), climate-friendly policies for the agricultural sector will produce greater impacts if they are based on the control and efficient use of local water resources.

Table 5 shows the mean sensitivities for the Brazilian Northeast and the states of the region. It is verified that, on average, the increase of 1 mm in the level of annual rainfall increases the value of agricultural production in the Northeast by 0.1874%, while the increase of 1 °c in the average annual temperature raises agricultural production by 0.2129% in the region.

Furthermore, it is possible to observe that Ceará was the state that showed the greatest average sensitivity in relation to changes in annual precipitation, followed by the states of Rio Grande do Norte and Paraíba. In relation to temperature, the highest average sensitivity was obtained by the state of Maranhão, followed by the states of Piauí and Ceará.

Considering the averages, only the states of Bahia and Maranhão show sensitivity levels to rainfall changes lower than the values obtained by the Northeast region as a whole. Regarding temperature, this behavior was observed for the states of Alagoas, Bahia, Ceará, Paraíba, Pernambuco, Rio Grande do Norte and Sergipe.

Table 5: Average State Sensitivity.

Estate	Precipitation	Temperature
Northeast	0.1804	0.2129
Alagoas	0.2080	0.1937
Bahia	0.1814	0.2063
Ceará	0.2129	0.2120
Maranhão	0.1330	0.2382
Paraíba	0.2098	0.2102
Pernambuco	0.2061	0.2041
Piauí	0.1975	0.2208
Rio Grande do Norte	0.2104	0.2070
Sergipe	0.1890	0.1963

Source: Elaborated by the authors.

Table 6 features a more detailed description of the average sensitivities of precipitation and temperature, presenting values per year for each Brazilian state in the region. Regarding the average sensitivities for precipitation, Table 6 shows that all the states in the region presented sensitivity growth in the year 2016, when compared to the initial year of the sample. It should be noted that the sensitivity of the agricultural sector to rainfall changes has already been verified for other localities, as discussed by [Kunwar e Bohara \(2017\)](#), which analyzed Nepal, and [Baylis, Paulson e Piras \(2011\)](#), which studied the United States of America.

Specifically, between 2006 and 2009, the state of Pernambuco was more sensitive to rainfall changes. In 2011, this characteristic was observed for the state of Sergipe, and from 2010 onwards, the greatest sensitivities to rainfall changes were observed in the state of Rio Grande do Norte.

Regarding the average sensitivity to shifts in temperature, it is verified that all the states in the region had their sensitivities decreased, when comparing the values of 2016 with the initial year of sample.

The greatest sensitivities were obtained by the State of Maranhão, from 2006 to 2016. This result is consistent with expectations, considering the geographic location of the state to the Northwest of the region, and its proximity to the Northern region of Brazil. Furthermore, according to [INPE \(2015\)](#), the annual rainfall level of Maranhão makes it stand out, presenting a tendency that differentiates it from other states regarding temperature effects.

Table 6: Average Sensitivity per year and state.

	Precipitation								
	AL	BA	CE	MA	PB	PE	PI	RN	SE
2006	0.2008	0.1503	0.1200	0.1037	0.1923	0.2051	0.1536	0.0716	0.1204
2007	0.1507	0.1646	0.1650	0.1254	0.2082	0.2091	0.1967	0.1283	0.0377
2008	0.1612	0.1637	0.1399	0.0876	0.1628	0.1557	0.1576	0.1253	0.0958
2009	0.1591	0.1600	0.1325	0.0786	0.1512	0.1736	0.1370	0.1034	0.1208
2010	0.1881	0.1665	0.2407	0.1390	0.2094	0.1844	0.2001	0.2544	0.1967
2011	0.2022	0.1748	0.1720	0.0866	0.1635	0.1781	0.1722	0.1866	0.2227
2012	0.2498	0.2082	0.2775	0.1822	0.2660	0.2453	0.2393	0.3043	0.2578
2013	0.2320	0.1960	0.2693	0.1419	0.2336	0.2193	0.2082	0.2878	0.2475
2014	0.2299	0.1890	0.2551	0.1307	0.2291	0.2103	0.2106	0.2722	0.2436
2015	0.2596	0.2109	0.2737	0.1809	0.2521	0.2334	0.2437	0.2936	0.2710
2016	0.2542	0.2117	0.2965	0.2069	0.2394	0.2330	0.2538	0.2872	0.2650
	Temperature								
2006	0.1916	0.2186	0.2203	0.2483	0.2097	0.2045	0.2331	0.2079	0.1901
2007	0.1890	0.2064	0.2103	0.2411	0.2057	0.1967	0.2159	0.2044	0.1974
2008	0.2057	0.2084	0.2323	0.2516	0.2288	0.2204	0.2303	0.2281	0.1982
2009	0.2160	0.2227	0.2424	0.2538	0.2322	0.2256	0.2358	0.2387	0.2204
2010	0.2178	0.2218	0.2097	0.2367	0.2159	0.2225	0.2236	0.2054	0.2147
2011	0.2043	0.2064	0.2298	0.2357	0.2329	0.2194	0.2269	0.2291	0.1950
2012	0.1669	0.1877	0.1910	0.2296	0.1832	0.1756	0.2074	0.1823	0.1695
2013	0.1803	0.2073	0.1983	0.2350	0.2037	0.1991	0.2211	0.1935	0.1864
2014	0.1850	0.1999	0.2040	0.2355	0.1999	0.1943	0.2155	0.1997	0.1834
2015	0.1824	0.1962	0.2014	0.2296	0.1955	0.1929	0.2109	0.1932	0.1788
2016	0.1818	0.1935	0.1923	0.2229	0.2045	0.1943	0.2083	0.1950	0.1814

Source: Elaborated by the authors.

In general, the results presented in Table 6 indicate that agricultural production in the Northeast region has become increasingly sensitive to variations in the precipitation levels, while changes in average temperatures have influenced to a lesser extent the value of agricultural production.

Historically, agricultural production in the Northeast region of Brazil has suffered from a lack of policies aimed at solving the problems arising from the region's water shortages, such as the application of measures to encourage the efficient use of water resources in farming. Also, these results indicate that public policies to combat the effects of climate change on agriculture in the Northeast region of Brazil will have greater impacts when based on the water needs of this region's municipalities.

Thus, it can be said that agriculture in the municipalities of the Northeast region of Brazil has suffered from a greater lack of policies aimed at solving the problems stemming from the water scarcity of the region. Also, the need to apply incentive measures to the efficient use of water resources in agriculture can be highlighted.

5 CONCLUDING REMARKS

The present work had as its objective to verify the sensibility of the agricultural production in the Northeast region of Brazil in relation to changes in precipitation and temperature levels. For that purpose, climate data was used, organized in terms of time period and geography, and the sensitivities were estimated by means of an econometric procedure for spatial panel data. i.e., the adopted econometric procedure includes the effects of the spatial proximity of the data units. Mean

precipitation and temperature data were used for a set of municipalities in the Northeast region of Brazil between 2006 and 2016.

The results showed that the agricultural production of the municipalities of the Brazilian Northeast is spatially autocorrelated over the years analyzed. The existence of spatial agglomerations of agricultural production in the Northeastern municipalities was verified through the calculation of the Moran Global Index, indicating that there are municipalities with high (or low) levels of production, which have neighbors with these same characteristics. Through the estimation of a spatial lag model, it was observed that changes in the production of a given municipality overflow spatially, also impacting the neighboring municipalities. Thus, public policies to support the agricultural sector of the Northeastern municipalities should be applied at a regional, and not just municipal, level, allowing greater efficiency.

With the calculation of the sensitivities, it was also verified that from 2006 to 2011, the agricultural production of the Brazilian Northeast was more sensitive to changes in temperature levels than to changes in average precipitation. However, after the years 2012, 2013, 2015 and 2016, agricultural production has become more sensitive to changes in the precipitation levels.

It was also noted that in the analyzed period there was an increase in the average sensitivity of the agricultural production in relation to the precipitation levels, while the average temperature sensitivity showed a decrease. In addition, it was verified that, on average, the agricultural production of the states of Pernambuco, Sergipe and Rio Grande do Norte present the greatest sensitivities to rainfall changes, while agriculture in the state of Maranhão is the most sensitive to changes in temperature levels.

Lastly, it can be concluded that the agricultural production in the Northeast region of Brazil has featured a lack of policies aimed at solving the problems arising from the region's water shortages, such as the application of measures to encourage the efficient use of water resources in farming. Furthermore, this work surmises that public policies to combat the effects of climate change on agriculture in the Northeast region of Brazil will have greater impacts when based on the water needs of this region's municipalities.

References

- AGRITEMPO. 2017. <<https://www.agritempo.gov.br/agritempo/index.jsp>>. Acessado em 17/12/2017.
- ALMEIDA, E. Econometria espacial aplicada. *Campinas-SP. Alínea*, 2012.
- ANSELIN, L. Interactive techniques and exploratory spatial data analysis. *Geographical Information Systems: principles, techniques, management and applications*, Citeseer, v. 1, p. 251–264, 1999.
- ARAÚJO, P. H. C. et al. Uma análise do impacto das mudanças climáticas na produtividade agrícola da região nordeste do brasil. *Revista Econômica do Nordeste*, v. 45, n. 3, p. 46–57, 2014.
- BARRETO, R. C. S.; ALMEIDA, E. S.; LIMA, J. E. de. Convergência espacial do pib per capita no estado do ceará. *Revista de Economia*, v. 36, n. 3, 2010.
- BAUMONT, C. *Spatial effects in housing price models: do housing prices capitalize urban development policies in the agglomeration of Dijon (1999)?* Tese (Doutorado) — Laboratoire d'économie et de gestion (LEG), 2004.
- BAYLIS, K.; PAULSON, N. D.; PIRAS, G. Spatial approaches to panel data in agricultural economics: a climate change application. *Journal of Agricultural and Applied Economics*, Cambridge University Press, v. 43, n. 3, p. 325–338, 2011.

- BURKE, M.; EMERICK, K. Adaptation to climate change: Evidence from us agriculture. *American Economic Journal: Economic Policy*, v. 8, n. 3, p. 106–40, 2016.
- CALZADILLA, A. et al. Climate change impacts on global agriculture. *Climatic change*, Springer, v. 120, n. 1-2, p. 357–374, 2013.
- CUNHA, D. A. d. et al. Irrigação como estratégia de adaptação de pequenos agricultores às mudanças climáticas: aspectos econômicos. *Revista de Economia e Sociologia Rural*, SciELO Brasil, v. 51, n. 2, p. 369–386, 2013.
- DOMINGUES, E. P.; MAGALHÃES, A. S.; RUIZ, R. M. Cenários de mudanças climáticas e agricultura no brasil: impactos econômicos na região nordeste. *Revista Econômica do Nordeste*, v. 42, n. 2, p. 229–246, 2011.
- ELHORST, J. P. Specification and estimation of spatial panel data models. *International regional science review*, Sage Publications, v. 26, n. 3, p. 244–268, 2003.
- ELHORST, J. P. Applied spatial econometrics: raising the bar. *Spatial Economic Analysis*, Taylor & Francis, v. 5, n. 1, p. 9–28, 2010.
- EVANGELISTA, P.; YOUNG, N.; BURNETT, J. How will climate change spatially affect agriculture production in ethiopia? case studies of important cereal crops. *Climatic change*, Springer, v. 119, n. 3-4, p. 855–873, 2013.
- FILHO, H. M. et al. Mudança do clima e os impactos na agricultura familiar no norte e nordeste do brasil. *Embrapa Solos-Artigo em periódico indexado (ALICE)*, IPC-IG Working Paper, Brasília, DF, n. 141, May 2016., 2016.
- FISCHER, G.; SHAH, M. M.; VELTHUIZEN, H. V. Climate change and agricultural vulnerability. IIASA, Laxenburg, Austria, 2002.
- IBGE. 2017. <https://www.ibge.gov.br/>.
- INPE. Cenários de mudanças climáticas: Regionalização. 2015.
- JU, H. et al. The impacts of climate change on agricultural production systems in china. *Climatic Change*, Springer, v. 120, n. 1-2, p. 313–324, 2013.
- KUNWAR, S. B.; BOHARA, A. K. Climate sensitivities and farmland values in nepal: A spatial panel ricardian approach. *Journal of Development and Agricultural Economics*, v. 9, n. 6, p. 145–161, 2017.
- LESAGE, J.; PACE, R. K. *Introduction to spatial econometrics*. [S.l.]: Chapman and Hall/CRC, 2009.
- MMA. 2017. <http://www.mma.gov.br/clima/convencao-das-nacoes-unidas/acordo-de-paris>). Acessado em 17/12/2017.
- MORAES, G. I. de; FILHO, J. B. de S. F. Brasil, mudanças climáticas e economia: o que há estabelecido? *Planejamento e Políticas Públicas*, n. 41, 2013.
- NASA. 2017. https://data.giss.nasa.gov/gistemp/graphs_v3/). Acessado em 17/12/2017.
- NELSON, G. C. et al. *Climate change: Impact on agriculture and costs of adaptation*. [S.l.]: Intl Food Policy Res Inst, 2009. v. 21.

- NELSON, G. C. et al. *Climate change: Impact on agriculture and costs of adaptation*. [S.l.]: Intl Food Policy Res Inst, 2009. v. 21.
- NYS, E. D.; ENGLE, N.; MAGALHÃES, A. Secas no brasil: política e gestão proativas. *Brasília, DF: Centro de Gestão e Estudos Estratégicos-CGEE*, 2016.
- OBERMAIER, M.; ROSA, L. P. Mudança climática e adaptação no brasil: uma análise crítica. *Estudos avançados*, SciELO Brasil, v. 27, n. 78, p. 155–176, 2013.
- OCHIENG, J.; KIRIMI, L.; MATHENGE, M. Effects of climate variability and change on agricultural production: The case of small scale farmers in kenya. *NJAS-Wageningen Journal of Life Sciences*, Elsevier, v. 77, p. 71–78, 2016.
- PAM. 2017. <https://sidra.ibge.gov.br/pesquisa/pam/tabelas>.
- PARRY, M. L. et al. Effects of climate change on global food production under sres emissions and socio-economic scenarios. *Global environmental change*, Elsevier, v. 14, n. 1, p. 53–67, 2004.
- PIAO, S. et al. The impacts of climate change on water resources and agriculture in china. *Nature*, Nature Publishing Group, v. 467, n. 7311, p. 43, 2010.
- SIDRA. 2017. <https://sidra.ibge.gov.br/home/ipca/brasil>.
- WIEBE, K. et al. Climate change impacts on agriculture in 2050 under a range of plausible socioeconomic and emissions scenarios. *Environmental Research Letters*, IOP Publishing, v. 10, n. 8, p. 085010, 2015.
- YALEW, A. W. et al. *Economic effects of climate change in developing countries: Economy-wide and regional analysis for Ethiopia*. [S.l.], 2017.